ATTACHMENT 6 **REVISED 02/2022** 

COMPANY: FORM(S) NUMBER: SERFF TRACKING NUMBER:		
MARKET TYPE	☐ Individual ☐ Small Group ☐ Large Group	
		1

Benefit	Description	Please specify location (Form/Page/Paragraph/ Other) of complying provision/language or attach explanation for an N/A response	FOR OFFICIAL USE ONLY
	Essential Health Benefits		
Air ambulance	Covered, No limits. Out of area air ambulance coverage is not covered.		
Allergy tests	(50) Test per policy year per enrollee <sup>1</sup> . Vaccines not covered.		
Bariatric Surgery	Subject to preauthorization. It must be covered the payment of (1) of the types of the bariatric surgery per member for life in Puerto Rico, if the services are available. The types of bariatric surgery that may be covered are the following: gastric bypass, adjustable band or sleeve gastrectomy. Coverage is available only to a diagnostic of morbid obesity. Morbid obesity means is the excess fat in the body determined by a body mass index (BMI) greater or equal to 35. The insured and dependents may have to meet a waiting period of 12 months before the benefit is covered, unless the physician certifies that the patient's life is in imminent danger. The facility must have accreditation form the Joint Commission and one of two entities the American College of Surgeon or the American Society for Metabolic and Bariatric Surgery. Surgeries to remove excess skin (commonly known as flaps) are not covered, unless the physician certifies that it is necessary to remove excess skin, since it affects the functionally of a limb or body part.		
Biophysical profile	(1) Procedure per pregnancy.		
Chiropractic Care	(20) Physical therapies or manipulations covered under a combined limit per policy year per enrollee.		
Contraceptive methods	Covered, No limits.		
Cryo-surgery of the uterus	(1) procedures per year per enrollee.		
Delivery and All Inpatient Services for Maternity Care  Dental Check-Up for Children	Delivery of baby (48) hour minimum length for vaginal delivery and (96) for cesarean delivery.  Covered for main enrollee, spouse and dependent daughter.  Dental checkup and cleanings (2) per policy year per member (every (6) months); bitewings and periapicals no more than one set every 3 years.		

<sup>\*</sup> Page number range are not accepted (i.e. 28-32)

\*\* The absence of explanation may prevent the filing approval.

<sup>&</sup>lt;sup>1</sup> Term use in this attachment to refer to an insured or subscriber

Benefit	Description	Please specify location (Form/Page/Paragraph/ Other) of complying provision/language or attach explanation for an N/A response	FOR OFFICIAL USE ONLY
Diagnostic Test (X-Ray and Lab Work)	No Limits. The reproduction of X-Rays must be covered. Laboratories related to infertility problems are covered as long as the same are laboratories covered in the policy.		
Dialysis and hemodialysis	(90) Days. Services related to any type of dialysis or hemodialysis, as well as services for any complication that may arise and their corresponding hospital or medical-surgical services. Will be covered for the first (90) days from: a) the date in which the member became eligible for the policy during the first time or, b) the date in which he/she received the first dialysis and hemodialysis. This will apply when subsequent dialysis or hemodialysis are related to the same clinical conditions.  These services must be covered for inpatient and outpatient services.		
Durable Medical Equipment	Covered with a preauthorization from plan rental or purchase or oxygen and necessary equipment for its administration/wheelchair/hospital bed. Mechanical respirators and ventilators are covered without limits as required by Law No. 62 of May 4, 2015 to member's patients under age of (21) and those who have started treatment as minors and meet (21) years and who received medical services or receive home care will continue to receive these services after (21) years of age. Coverage include also the following benefits: technological equipment necessary to enable the insured to stay alive; at least one daily shift of (8) hours of skilled nursing services with expertise in respiratory therapy or respiratory therapy specialists with expertise in nursing; supply that involve the management of the technological equipment; physical and occupational therapy.		
Emergency Room Services	No Limits. No preauthorization or waiting period required. Emergency services for outnetwork providers cannot be covered through reimbursement.  Limitations indicating that the emergency service must be received during the first 24 hours are not accepted.  Carrier with emergency telephone lines that offers waiver or a lower copayment or coinsurance if the member calls to such line cannot make any difference between an innetwork or out-network provider.		
Emergency Transportation/Ambulance	Services requested through the 9-1-1 Emergency System, covered and paid directly to the provider.  Other transportation services (i.e. transportation between institutions) covered by reimbursement up to \$80 per trip.		

<sup>\*</sup> Page number range are not accepted (i.e. 28-32)

\*\* The absence of explanation may prevent the filing approval.

<sup>&</sup>lt;sup>1</sup> Term use in this attachment to refer to an insured or subscriber

Benefit	Description	Please specify location (Form/Page/Paragraph/ Other) of complying provision/language or attach explanation for an N/A response	FOR OFFICIAL USE ONLY
Eye Glasses for Children	(1) pair of glasses (lenses and frames per year per member), supplemented using FEDVIP. Low Vision coverage - Glasses for member until age (21). (1) pair per policy year per member within the contracted collection, including high-powered glasses to policyholders with significant loss of vision, but do not have totally blind. Also covers one item per year per member, visual aids (prescription lenses, telescopes single or double lens) to policyholders up to age (21) with significant loss of vision, but do not have totally blind. Services related to Eye glasses cannot be provided through reimbursement, discounts or allowance.		
Gastrointestinal endoscopies	Covered, No Limits.		
Generic Drugs	Subject to a Drug List, Generics as a first option. Some medications may require preauthorization, age limits, quantity limits, specialty limits and/or step therapy. Drugs related to mental health conditions cannot include limitations indicating that a specific drug must be prescribe by a psychiatrist or neurologist.  Not all the drugs related to mental/behavioral health conditions can be manage through a specialty limit.		
Habilitation Services	(20) Physical therapies or manipulations covered under a combined limit per year. Services limited to physical therapies, except for those covered under home health care benefit.		
Home Health Care Services	(40) physical, occupational and speech therapy under a combined limit per policy year per member. Covered only if they begin 14 days after members discharge from hospital of at least (3) days and if they are provided for the same condition by he/she was admitted.		
Imaging (CT/PET Scans, MRIs)	For PET & PET/CT, (1) Per policy year per enrollee. For MRI & CT, (1) per anatomical region per policy year per enrollee.		
Inpatient Hospital Services (e.g., Hospital Stay)	Excludes services for personal comfort and or custodial services. Hospitalizations for services or procedures that may be performed in an outpatient services are not covered.		
Inpatient Physician and Surgical Services	Covered, No Limits.		
Intra-articular injections	(12) Injections per policy year per member, up to (2) daily injections.		
procedures and tests	Electromyograms covered up to (2) procedures per year per enrollee.		
Lithotripsy	Covered, No limits		

<sup>\*</sup> Page number range are not accepted (i.e. 28-32)
\*\* The absence of explanation may prevent the filing approval.

<sup>&</sup>lt;sup>1</sup> Term use in this attachment to refer to an insured or subscriber

Benefit	Description	Please specify location (Form/Page/Paragraph/ Other) of complying provision/language or attach explanation for an N/A response	FOR OFFICIAL USE ONLY
Mental/Behavioral Health Inpatient Services	Residential treatment outside service area is not covered. No limit in accordance to the Mental Health Parity Act. Limitations indicating that these services must be received or coordinate through an specific company or program are not accepted.		
Mental/Behavioral Health Outpatient Services	No limit in accordance to the Mental Health Parity Act. Limitations indicating that these services must be received or coordinate through an specific company or program are not accepted.		
MRA	Covered, No limits.		
Nerve conduction velocity tests	(2) Procedures per policy year per enrollee.		
Neurological tests and procedures	Covered, No limits.		
Non-Preferred Brand Drugs	Subject to a Drug List, Generics as a first option. Some medications may require preauthorization, age limits, quantity limits, specialty limits and/or step therapy. Drugs related to mental health conditions cannot include limitations indicating that a specific drug must be prescribe by a psychiatrist or neurologist.  Not all the drugs related to mental/behavioral health conditions can be manage through a		
	specialty limit.		
Nuclear medicine tests	Covered, No Limits.		
Nutritionist services	(4) Per policy year per enrollee. Limited to morbid, renal and diabetes conditions. Covered by reimbursement up to \$20 per visit.		
Oral, Intravenously, Injectable or Intrathecal chemotherapy	Covered, No Limits. Antineoplastic agents cannot be excluded from the basic coverage.		
	These services must be covered for inpatient and outpatient services.		
Orthognatic surgery	Expenses related for materials are excluded.		
Other Practitioner Office Visit (Nurse, Physician Assistant)	Non physician professionals or doctors in odontology including nurse and physician assistant except those required by local law such as: podiatrist, audiologist, optometrist, clinical psychologists and chiropractors.		
Out of area coverage (US)	Services are covered for emergency cases or cases that required equipment, treatment and facilities not available in Puerto Rico. Services are subject to preauthorization from the plan except for an emergency. Elective treatments, not considered as an emergency, are not covered by this policy. Rates to be paid are the usual and customary (UCR) rate of the geographical area in which the services are provided.		
Outpatient Facility Fee (e.g., Ambulatory Surgery Center)	Services rendered in an outpatient facility that may be performed in physician's office are not covered.		

<sup>\*</sup> Page number range are not accepted (i.e. 28-32)
\*\* The absence of explanation may prevent the filing approval.

<sup>&</sup>lt;sup>1</sup> Term use in this attachment to refer to an insured or subscriber

Benefit	Description	Please specify location (Form/Page/Paragraph/ Other) of complying provision/language or attach explanation for an N/A response	FOR OFFICIAL USE ONLY
Outpatient Rehabilitation Services	(20) Physical therapies or manipulations covered under a combined limit per policy year per enrollee.  Services not covered include occupational, speech and language therapies, prosthetics and implants, orthopedics and orthotic devices, cardiac rehabilitation.  Occupational, speech and language therapies must be covered for autism condition and for home health care services.  Services limited to physical therapies, except for those covered under home health care benefit.		
Outpatient Surgery Physician/Surgical Services	Excludes: Cosmetic surgery, oral surgery that is dental in origin except those as a result of an accident, mammoplasty (except those required for patients after a breast cancer mastectomy), septoplasty, blepharoplasty, rinoseptoplasty, procedures to re- establish the ability to procreate, organ transplant procedures, other than the specified in the transplant services benefit (other organ transplant may be covered as an optional benefit), induced abortion, experimental procedures, skin tags removal, ptosis repair, nail excisions, scalenotomy, Lasik and other surgical procedures to correct refractive defects, surgical assistance services, intravenous analgesia services or analgesia administered though inhalation at the physician or dentist's office, services for the treatment of the temporamandibular articulation syndrome, excision of granulomas or radicular cysts originated by infection in the tooth pulp; services to correct the vertical dimension or occlusion, removal of exostosis (mandibulary or maxillary).		
Polysomnography	(1) Type of test per lifetime per enrollee.		
Preferred Brand Drugs	Subject to a Drug List, Generics as a first option. Some medications may require preauthorization, age limits, quantity limits, specialty limits and/or step therapy. Drugs related to mental health conditions cannot include limitations indicating that a specific drug must be prescribe by a psychiatrist or neurologist.  Not all the drugs related to mental/behavioral health conditions can be manage through a specialty limit.		
Prenatal and Postnatal Care	Covered for mainholder, spouse and dependent daughter.		
Preventive Care/Screening/Immunization	Preventive care that meets recommendations described below, as established by the USPSTF.		
Primary Care Visit to Treat an Injury or Illness	Covered, No Limits.		
Radiation therapy	Covered, No Limits. These services must be covered for inpatient and outpatient services.		

<sup>\*</sup> Page number range are not accepted (i.e. 28-32)

\*\* The absence of explanation may prevent the filing approval.

<sup>&</sup>lt;sup>1</sup> Term use in this attachment to refer to an insured or subscriber

Benefit	Description	Please specify location (Form/Page/Paragraph/ Other) of complying provision/language or attach explanation for an N/A response	FOR OFFICIAL USE ONLY
Routine Dental Services (Adult)	Dental checkup and cleaning (2) per policy year per member (every (6) months); bitewings and periapicals no more that one set every (3) years. (Optional coverage; Orthodontic, Periodontics, Endodontic, prosthetic dental services, Full mouth reconstructions, Fluoride treatment covered to members under age (19) and Root canal only to anterior and posterior teeth.)		
Routine Eye Exam (Adult)	Refraction exam is covered (1) per year, per member.		
Routine Eye Exam for Children	(1) Visit per year supplemented using FEDVIP.		
Routine Foot Care	Covered, No Limits.		
Skilled Nursing Facility	Covered only if they begin (14) days after member's discharge from hospital of at least (3) days and if they are provided for the same condition by he/she was admitted. Maximum of (120) days.		
Specialist Visit	Covered, No Limits.		
Specialty Drugs	Subject to a Drug List, Generics as a first option. Some medications may require preauthorization, age limits, quantity limits, specialty limits and/or step therapy. Drugs related to mental health conditions cannot include limitations indicating that a specific drug must be prescribe by a psychiatrist or neurologist.  Not all the drugs related to mental/behavioral health conditions can be manage through a specialty limit.		
Sterilization	Covered, No Limits.		
Substance Abuse Disorder Inpatient Services	No limits in accordance to the Mental Health Parity Act. Include Detox and Residential treatment services. Residential treatment outside service area is not covered.  Partials are included: (2) partial hospital days equivalent to (1) regular day.  Limitations indicating that these services must be received or coordinate through an specific company or program are not accepted.  Expenses for services resulting from the administration of an employer drug detection program are not covered. However, after the enrollee's participation in any treatment related to a positive outcome in the employer drug detention program, he/she is eligible for treatment under the contract.  Language indicating that the emergency services must be received during the first 24 hours is not accepted.  Not all the drugs related to mental/behavioral health conditions can be manage through a specialty limit.		

<sup>\*</sup> Page number range are not accepted (i.e. 28-32)
\*\* The absence of explanation may prevent the filing approval.

<sup>&</sup>lt;sup>1</sup> Term use in this attachment to refer to an insured or subscriber

Benefit	Description	Please specify location (Form/Page/Paragraph/ Other) of complying provision/language or attach explanation for an N/A response	FOR OFFICIAL USE ONLY
Substance Abuse Disorder Outpatient Services	No limits in accordance to the Mental Health Parity Act.  Limitations indicating that these services must be received or coordinate through an specific company or program are not accepted.  Expenses for services resulting from the administration of an employer drug detection program are not covered. However, after the enrollee's participation in any treatment related to a positive outcome in the employer drug detention program, he/she is eligible for treatment under the contract.  Language indicating that the emergency services must be received during the first 24 hours is not accepted.  Not all the drugs related to mental/behavioral health conditions can be manage through a specialty limit.		
Transplant Services	Benefit covers skin, bone and comeal transplants. Coverage extends to charges directly related to the transplant service, including care prior to surgery, post-surgery care and treatment in respect of immunosuppressive drugs.  Services are covered only through participating providers. Pre authorization is required. This benefit will be covered 100% in Puerto Rico and should not be available through reimbursement.		
Tympanometry	(1) Per policy year per enrollee.		
Urgent Care Centers or Facilities	No Limits. No preauthorization or waiting period required.		

<sup>\*</sup> Page number range are not accepted (i.e. 28-32)

\*\* The absence of explanation may prevent the filing approval.

<sup>&</sup>lt;sup>1</sup> Term use in this attachment to refer to an insured or subscriber

Benefit	Description	Please specify location (Form/Page/Paragraph/ Other) of complying provision/language or attach explanation for an N/A response	FOR OFFICIAL USE ONLY	
**ADULTS PREVENTIVE S	Covered Preventive Services for Adult (NO COST SHARING IS APPLICABLE) ERVICES MUST BE INCLUDED IN THE SAME ORDER AND DESCRIPTION ESTAB	BLISHED IN THIS CHECKLIST	**	
Abdominal Aortic Aneurysm	(1) time screening for abdominal aortic aneurysm (AAA) by ultrasonography in men aged (65) to (75) who have ever smoked.			
Aspirin Use to Prevent Cardiovascular Disease and Colorectal Cancer: Preventive Medication	The USPSTF recommends initiating low-dose aspirin use for the primary prevention of cardiovascular disease and colorectal cancer in adults aged 50 to 59 years who have a 10% or greater 10-year cardiovascular risk, are not at increased risk for bleeding, have a life expectancy of at least 10 years, and are willing to take low-dose aspirin daily for at least 10 years.			
Colorectal Cancer	The USPSTF recommends screening for colorectal cancer (CRC) using fecal occult blood testing, sigmoidoscopy, or colonoscopy, in adults, beginning at age (45) years and continuing until age (75) years. The risks and benefits of these screening methods vary.		U	J <b>pdated</b>
Depression screening for Adults	The USPSTF recommends screening for depression in the general adult population, including pregnant and postpartum women. Screening should be implemented with adequate systems in place to ensure accurate diagnosis, effective treatment, and appropriate follow-up.			
Falls prevention in older adults: Vitamin D	The USPSTF recommends vitamin D supplementation to prevent falls in community-dwelling adults age 65 years and older who are at increased risk for falls.			
Healthy diet and physical activity counseling to prevent cardiovascular disease: adults with cardiovascular risk factors	The USPSTF recommends offering or referring adults who are overweight or obese and have additional cardiovascular disease (CVD) risk factors to intensive behavioral counseling interventions to promote a healthful diet and physical activity for CVD prevention.			
Hepatitis B screening: Nonpregnant Adolescents and Adults	The USPSTF recommends screening for hepatitis B virus (HBV) infection in persons at high risk for infection.			
Hepatitis C virus infection screening: Adolescents and Adults	The USPSTF recommends screening for hepatitis C virus (HCV) infection in adults aged 18 to 79 years.			

<sup>\*</sup> Page number range are not accepted (i.e. 28-32)
\*\* The absence of explanation may prevent the filing approval.

<sup>&</sup>lt;sup>1</sup> Term use in this attachment to refer to an insured or subscriber

Benefit	Description	Please specify location (Form/Page/Paragraph/ Other) of complying provision/language or attach explanation for an N/A response	FOR OFFICIAL USE ONLY	
Hypertension Screening in Adults 18 years or older without known hypertension	The USPSTF recommends screening for hypertension in adults 18 years or older with office blood pressure measurement. The USPSTF recommends obtaining blood pressure measurements outside of the clinical setting for diagnostic confirmation before starting treatment.			Updated
HIV preexposure prophylaxis for the prevention of HIV infection	The USPSTF recommends that clinicians offer preexposure prophylaxis (PrEP) with effective antiretroviral therapy to persons who are at high risk of HIV acquisition.			
HIV screening: nonpregnant adolescents and adults	Clinicians screening for HIV infection in adolescents and adults ages (15) to (65) years. Younger adolescents and older adults who are at increased risk should also be screened. As required by Law 45-2016 one HIV test a year as part of the routine studies for any medical evaluation, except for pregnant women to which apply the requirements of the USPSTF. You should consider the recommendations set out in the Department of Health's HIV regulations.			
Immunization	Vaccines for adults-doses, recommended ages, and recommended populations vary:  Hepatitis A, Hepatitis B, Herpes Zoster, Human Papillomavirus, Influenza (Flu Shot),  Measles, Mumps, Rubella, Meningococcal, Pneumococcal, Tetanus, Diphtheria,  Pertussis, Varicella. Catch ups must be covered.  COVID-19 vaccine as part of preventive immunizations for adolescents over 16 years of age and adults, according to the order established by the Puerto Rico Department of Health and the recommendations of the Advisory Committee on Immunization Practices (ACIP), of the Centers for Disease Control and Prevention (CDC), in accordance with Article 2.050 (C) of the Health Insurance Code of Puerto Rico.			

<sup>\*</sup> Page number range are not accepted (i.e. 28-32)
\*\* The absence of explanation may prevent the filing approval.

<sup>&</sup>lt;sup>1</sup> Term use in this attachment to refer to an insured or subscriber

Benefit	Description	Please specify location (Form/Page/Paragraph/ Other) of complying provision/language or attach explanation for an N/A response	FOR OFFICIAL USE ONLY	
Lung cancer screening	The USPSTF recommends annual screening for lung cancer with low-dose computed tomography (LDCT) in adults ages (50) to (80) years who have a (20) pack-year smoking history and currently smoke or have quit within the past (15) years. Screening should be discontinued once a person has not smoked for (15) years or develops a health problem that substantially limits life expectancy or the ability or willingness to have curative lung surgery.			
Obesity screening and counseling: adults	The USPSTF recommends that clinicians offer or refer adults with a body mass index of 30 or higher (calculated as weight in kilograms divided by height in meters squared) to intensive, multicomponent behavioral interventions.			
Prediabetes and Type 2 Diabetes Screening in asymptomatic adults aged 35 to 70 years who have overweight or obesity	The USPSTF recommends screening for prediabetes and type 2 diabetes in adults aged 35 to 70 years who have overweight or obesity. Clinicians should offer or refer patients with prediabetes to effective preventive interventions.			Updated
Sexually Transmitted Infections (STI) Counseling	The USPSTF recommends intensive behavioral counseling for all sexually active adolescents and for adults who are at increased risk for sexually transmitted infections (STIs).			
Statin Use for the Primary Prevention of Cardiovascular Disease in Adults: Preventive Medication	The USPSTF recommends that adults without a history of cardiovascular disease (CVD) (i.e., symptomatic coronary artery disease or ischemic stroke) use a low- to moderate-dose statin for the prevention of CVD events and mortality when all of the following criteria are met: 1) they are ages 40 to 75 years; 2) they have 1 or more CVD risk factors (i.e., dyslipidemia, diabetes, hypertension, or smoking); and 3) they have a calculated 10-year risk of a cardiovascular event of 10% or greater. Identification of dyslipidemia and calculation of 10-year CVD event risk requires universal lipids screening in adults ages 40 to 75 years.			

<sup>\*</sup> Page number range are not accepted (i.e. 28-32)

\*\* The absence of explanation may prevent the filing approval.

<sup>&</sup>lt;sup>1</sup> Term use in this attachment to refer to an insured or subscriber

Benefit	Description	Please specify location (Form/Page/Paragraph/ Other) of complying provision/language or attach explanation for an N/A response	FOR OFFICIAL USE ONLY
Syphilis screening: nonpregnant persons and adolescents	The USPSTF recommends screening for syphilis infection in persons who are at increased risk for infection.		

<sup>\*</sup> Page number range are not accepted (i.e. 28-32)

\*\* The absence of explanation may prevent the filing approval.

<sup>&</sup>lt;sup>1</sup> Term use in this attachment to refer to an insured or subscriber

Benefit	Description	Please specify location (Form/Page/Paragraph/ Other) of complying provision/language or attach explanation for an N/A response	FOR OFFICIAL USE ONLY
Tobacco Smoking cessation and Medication: nonpregnant adults	The USPSTF recommends that clinicians ask all adults about tobacco use, advise them to stop using tobacco, and provide behavioral interventions and U.S. Food and Drug Administration (FDA)—approved pharmacotherapy for cessation to nonpregnant adults who use tobacco.  For those who uses tobacco cessation products, this plan cover the dispatch of FDA approved medication for smoke cessation for (90) consecutive days in one intent and until (2) intents per year. The recommendation does not established any difference on the medication for smoke cessation tier. If the formulary include the drug no cost sharing can be applied regardless the specific tier.		
Tuberculosis screening: adults	The USPSTF recommends screening for latent tuberculosis infection in populations at increased risk.		
Unhealthy alcohol use: adults	The USPSTF recommends screening for unhealthy alcohol use in primary care settings in adults 18 years or older, including pregnant women, and providing persons engaged in risky or hazardous drinking with brief behavioral counseling interventions to reduce unhealthy alcohol use.		
Unhealthy Drug Use: adults	The USPSTF recommends screening by asking questions about unhealthy drug use in adults age 18 years or older. Screening should be implemented when services for accurate diagnosis, effective treatment, and appropriate care can be offered or referred. (Screening refers to asking questions about unhealthy drug use, not testing biological specimens.)		

<sup>\*</sup> Page number range are not accepted (i.e. 28-32)

\*\* The absence of explanation may prevent the filing approval.

<sup>&</sup>lt;sup>1</sup> Term use in this attachment to refer to an insured or subscriber

Benefit	Description	Please specify location (Form/Page/Paragraph/ Other) of complying provision/language or attach explanation for an N/A response	FOR OFFICIAL USE ONLY
	ered Preventive Services for Women, Including Pregnant (NO COST SHARING IS APPLERVICES MUST BE INCLUDED IN THE SAME ORDER AND DESCRIPTION ESTAI	· · · · · · · · · · · · · · · · · · ·	<b>[**</b>
Asymptomatic Bacteriuria Screening: pregnant persons	The USPSTF recommends screening for asymptomatic bacteriuria using urine culture in pregnant persons.		
BRCA-Related Cancer: Risk Assessment, Genetic Counseling, and Genetic Testing	The USPSTF recommends that primary care clinicians assess women with a personal or family history of breast, ovarian, tubal, or peritoneal cancer or who have an ancestry associated with breast cancer susceptibility 1 and 2 (BRCA1/2) gene mutations with an appropriate brief familial risk assessment tool. Women with a positive result on the risk assessment tool should receive genetic counseling and, if indicated after counseling, genetic testing.		
Breast Cancer Preventive Medication Use to Reduce Risk	The USPSTF recommends that clinicians offer to prescribe risk-reducing medications, such as tamoxifen, raloxifene, or aromatase inhibitors, to women who are at increased risk for breast cancer aged 35 years or older and at low risk for adverse medication effects.		
Breast Cancer Screening	The USPSTF recommends screening mammography for women, with or without clinical breast examination, every 1 to 2 years for women age 40 years and older. The USPSTF recommends biennial screening mammography for women aged (50) to (74) years.		

<sup>\*</sup> Page number range are not accepted (i.e. 28-32)
\*\* The absence of explanation may prevent the filing approval.

<sup>&</sup>lt;sup>1</sup> Term use in this attachment to refer to an insured or subscriber

Benefit	Description	Please specify location (Form/Page/Paragraph/ Other) of complying provision/language or attach explanation for an N/A response	FOR OFFICIAL USE ONLY	
Breastfeeding	Comprehensive lactation support services (including consultation; counseling; education by clinicians and peer support services; and breastfeeding equipment and supplies) during the antenatal, perinatal, and postpartum periods to optimize the successful initiation and maintenance of breastfeeding.  Breastfeeding equipment and supplies include, but are not limited to, double electric breast pumps (including pump parts and maintenance) and breast milk storage supplies. Access to double electric pumps should be a priority to optimize breastfeeding and should not be predicated on prior failure of a manual pump. Breastfeeding equipment may also include equipment and supplies as clinically indicated to support dyads with breastfeeding difficulties and those who need additional services. No monetary limits apply.  Breastfeeding equipment and supplies cannot be provided through reimbursement or in the optional Major Medical coverage.			Updated
Cervical Cancer Screening	The USPSTF recommends screening for cervical cancer every 3 years with cervical cytology alone in women aged 21 to 29 years.  For women aged 30 to 65 years, the USPSTF recommends screening every 3 years with cervical cytology alone, every 5 years with high-risk human papillomavirus (hrHPV) testing alone, or every 5 years with hrHPV testing in combination with cytology (cotesting).			
Chlamydia Screening in all sexually active women, including pregnant persons	The USPSTF recommends screening for chlamydia in all sexually active women 24 years or younger and in women 25 years or older who are at increased risk for infection.			Updated

<sup>\*</sup> Page number range are not accepted (i.e. 28-32)

\*\* The absence of explanation may prevent the filing approval.

<sup>&</sup>lt;sup>1</sup> Term use in this attachment to refer to an insured or subscriber

Benefit	Description	Please specify location (Form/Page/Paragraph/ Other) of complying provision/language or attach explanation for an N/A response	FOR OFFICIAL USE ONLY	
Contraception	All Food and Drug Administration approved, granted or cleared contraceptive methods, sterilization procedures, screening, patient education, counseling and provision of contraceptives (including in the immediate postpartum period) for all women with reproductive capacity as prescribed. Contraceptives care also includes follow-up care.  The full range of contraceptives includes those currently listed in the FDA's Birth Control Guide and any additional contraceptives approved, granted or cleared by the FDA. Any device insertion and removal of contraceptive methods is covered. Contraceptives methods cannot be provided through reimbursement. The Health Insurance Code of Puerto Rico, neither the federal applicable law established any difference in contraceptive tiers. If the formulary include the contraceptive no cost sharing can be applied regardless the specific tier.			Updated
Counseling for Healthy Weight and Weight Gain in Pregnancy: Pregnant Persons	The USPSTF recommends that clinicians offer pregnant persons effective behavioral counseling interventions aimed at promoting healthy weight gain and preventing excess gestational weight gain in pregnancy.			New
Intimate Partner Violence, Elder Abuse, and Abuse of Vulnerable Adults: Screening: women of reproductive age	The USPSTF recommends that clinicians screen for intimate partner violence (IPV) in women of reproductive age and provide or refer women who screen positive to ongoing support services.			
Folic Acid	The USPSTF recommends that all women planning or capable of pregnancy take a daily supplement containing (0.4) to (0.8) mg (400 to 800pg) of folic acid.			
Gestational Diabetes: Screening: asymptomatic pregnant persons at 24 weeks of gestation or after	The USPSTF recommends screening for gestational diabetes in asymptomatic pregnant persons at 24 weeks of gestation or after.			Updated
Gonorrhea Screening in all sexually active women, including pregnant persons	The USPSTF recommends screening for gonorrhea in all sexually active women 24 years or younger and in women 25 years or older who are at increased risk for infection			Updated
Hepatitis B Screening: Pregnant women	Screening for pregnant women at their first prenatal visit.			

<sup>\*</sup> Page number range are not accepted (i.e. 28-32)

\*\* The absence of explanation may prevent the filing approval.

<sup>&</sup>lt;sup>1</sup> Term use in this attachment to refer to an insured or subscriber

Benefit	Description	Please specify location (Form/Page/Paragraph/ Other) of complying provision/language or attach explanation for an N/A response	FOR OFFICIAL USE ONLY	
Human Immunodeficiency Virus (HIV) screening: Pregnant persons	The USPSTF recommends that clinicians screen for HIV infection in all pregnant persons, including those who present in labor or at delivery whose HIV status is unknown. Regarding pregnant women, all insurers or health services organizations are required to cover and will not impose cost-sharing requirements with regard to the following tests included in the most recent recommendations of the USPSTF:  1) A first HIV test during the first trimester of pregnancy at the first prenatal visit, and 2) A second test during the third trimester of pregnancy (between the (28) and (34) weeks of pregnancy).			
Screening for Human Immunodeficiency Virus Infection (HIV)	All adolescent and adult women, ages 15 and older, receive a screening test for HIV at least once during their lifetime. Earlier or additional screening should be based on risk, and rescreening annually or more often may be appropriate beginning at age 13 for adolescent and adult women with an increased risk of HIV infection.  Risk assessment and prevention education for HIV infection beginning at age 13 and continuing as determined by risk.  A screening test for HIV is recommended for all pregnant women upon initiation of prenatal care with rescreening during pregnancy based on risk factors. Rapid HIV testing is recommended for pregnant women who present in active labor with an undocumented HIV status. Screening during pregnancy enables prevention of vertical transmission.			Updated
Obesity Prevention in Midlife Women	Counseling midlife women aged 40 to 60 years with normal or overweight body mass index (BMI) (18.5-29.9 kg/m2) to maintain weight or limit weight gain to prevent obesity. Counseling may include individualized discussion of healthy eating and physical activity.		1	New
Osteoporosis screening: postmenopausal women younger than 65 years at increased risk of osteoporosis	The USPSTF recommends screening for osteoporosis with bone measurement testing to prevent osteoporotic fractures in postmenopausal women younger than 65 years who are at increased risk of osteoporosis, as determined by a formal clinical risk assessment tool.			
Osteoporosis screening: women 65 years and older	The USPSTF recommends screening for osteoporosis with bone measurement testing to prevent osteoporotic fractures in women 65 years and older.			

<sup>\*</sup> Page number range are not accepted (i.e. 28-32)

\*\* The absence of explanation may prevent the filing approval.

<sup>&</sup>lt;sup>1</sup> Term use in this attachment to refer to an insured or subscriber

Benefit	Description	Please specify location (Form/Page/Paragraph/ Other) of complying provision/language or attach explanation for an N/A response	FOR OFFICIAL USE ONLY	
Perinatal depression: counseling and intervention	The USPSTF recommends that clinicians provide or refer pregnant and postpartum persons who are at increased risk of perinatal depression to counseling interventions.			
Preeclampsia prevention: aspirin in pregnant persons at high risk for preeclampsia	The USPSTF recommends the use of low-dose aspirin (81 mg/day) as preventive medication after 12 weeks of gestation in persons who are at high risk for preeclampsia.			Updated
Preeclampsia screening	The USPSTF recommends screening for preeclampsia in pregnant women with blood pressure measurements throughout pregnancy.			
RH Incompatibility screening	Rh (D) blood typing and antibody testing for all pregnant women during their first visit for pregnancy- related care.  Also, repeated Rh (D) antibody testing for all unsensitized Rh (D)- negative women at (24)-(28) weeks gestation, unless the biological father is known to be Rh (D)-negative.			
Screening for anxiety	The Women's Preventive Services Initiative recommends screening for anxiety in adolescent and adult women, including those who are pregnant or postpartum. Optimal screening intervals are unknown and clinical judgement should be used to determine screening frequency. Given the high prevalence of anxiety disorders, lack of recognition in clinical practice, and multiple problems associated with untreated anxiety, clinicians should consider screening women who have not been recently screened.			

<sup>\*</sup> Page number range are not accepted (i.e. 28-32)
\*\* The absence of explanation may prevent the filing approval.

<sup>&</sup>lt;sup>1</sup> Term use in this attachment to refer to an insured or subscriber

Benefit	Description	Please specify location (Form/Page/Paragraph/ Other) of complying provision/language or attach explanation for an N/A response	FOR OFFICIAL USE ONLY
Screening for diabetes mellitus after pregnancy	Screening for diabetes mellitus in women with a history of gestational diabetes mellitus (GDM) who are not currently pregnant and who have not previously been diagnosed with type 2 diabetes mellitus. Initial testing should ideally occur within the first year postpartum and can be conducted as early as 4–6 weeks postpartum.		
Syphilis screening: pregnant women	Early screening for syphilis infection in all pregnant women.		

<sup>\*</sup> Page number range are not accepted (i.e. 28-32)
\*\* The absence of explanation may prevent the filing approval.

<sup>&</sup>lt;sup>1</sup> Term use in this attachment to refer to an insured or subscriber

Benefit	Description	Please specify location (Form/Page/Paragraph/ Other) of complying provision/language or attach explanation for an N/A response	FOR OFFICIAL USE ONLY
Screening for urinary incontinence in women	The Women's Preventive Services Initiative recommends screening women for urinary incontinence as a preventive service. Factors associated with an increased risk for urinary incontinence include increasing parity, advancing age, and obesity; however, these factors should not be used to limit screening.  Several screening tools demonstrate fair to high accuracy in identifying urinary incontinence in women. Although minimum screening intervals are unknown, given the prevalence of urinary incontinence, the fact that many women do not volunteer symptoms, and the multiple, frequently-changing risk factors associated with incontinence, it is reasonable to conduct annually.		
Tobacco Smoking cessation: Pregnant Persons	The USPSTF recommends that clinicians ask all pregnant persons about tobacco use, advise them to stop using tobacco, and provide behavioral interventions for cessation to pregnant persons who use tobacco.		
Well-woman visits	Well-woman preventive care visit annually (depending on the woman's health status, health needs an other risk factors) for adult women to obtain the recommended preventive services that are age and developmentally appropriate, including preconception care and many services necessary for prenatal care and interpregnnacy visits. This well-woman visit should, where appropriate, include other preventive services listed. If the clinician determines that a patient requires additional well-woman visits, the additional visits must be provided without cost sharing. The full range of contraceptives includes those currently listed in the Food and Drug Administration (FDA's) Birth Control Guide- approved,-granted, or-cleared contraceptives, effective family planning practices, and sterilization procedures be available as part of contraceptive care.		U
**CHILDREN PREVENTIV	Covered Preventive Services for Children (NO COST SHARING IS APPLICABLE)  E SERVICES MUST BE INCLUDED IN THE SAME ORDER AND DESCRIPTION ESTA		ST**
Anemia/Iron	Perform risk assessment or screening as appropriate, per recommendations in the current edition of the AAP Pediatric Nutrition: Policy of the American Academy of Pediatrics (Iron chapter).		

<sup>\*</sup> Page number range are not accepted (i.e. 28-32)

\*\* The absence of explanation may prevent the filing approval.

<sup>&</sup>lt;sup>1</sup> Term use in this attachment to refer to an insured or subscriber

Benefit	Description	Please specify location (Form/Page/Paragraph/ Other) of complying provision/language or attach explanation for an N/A response	FOR OFFICIAL USE ONLY
Autism Spectrum Disorder Screening	Screening for children at (18) and (24) months.		
Behavioral	Assessment for children of all ages. Ages: (0) to (9) months, (1) to (4) years, (5) to (10) years, (11) to (21) years.		
Bilirubin concentration	Screening for newborns		
Blood Pressure	Screening for Children Ages: (0) to (11) months, (1) to (4) years, (5) to (10) years, (11) to (14) years, (15) to (17) years.		
Blood screening	Screening for newborns		
Cervical Dysplasia	Screening for sexually active females.		
Depression screening: adolescents	The USPSTF recommends screening for major depressive disorder (MDD) in adolescents aged 12 to 18 years. Screening should be implemented with adequate systems in place to ensure accurate diagnosis, effective treatment, and appropriate follow-up.		
Developmental Screening and Surveillance	Screening for children under age (3), and surveillance throughout childhood.		
Dyslipidemia	Screening for all children once between 9 and 11 years and once between 17 and 21 years, and for children at higher risk of lipid disorders ages: 1 to 4 years, 5 to 10 years, 11 to 14 years, 15 to 17 years		
Gonorrhea prophylactic medication: newborns	The USPSTF recommends prophylactic ocular topical medication for all newborns to prevent gonococcal ophthalmia neonatorum.		
Hearing	Screening for hearing loss all newborns infants and for children once between 11 and 14 years, once between 15 and 17 years, and once between 18 and 21 years.		

<sup>\*</sup> Page number range are not accepted (i.e. 28-32)

\*\* The absence of explanation may prevent the filing approval.

<sup>&</sup>lt;sup>1</sup> Term use in this attachment to refer to an insured or subscriber

Benefit	Description	Please specify location (Form/Page/Paragraph/ Other) of complying provision/language or_attach explanation for an N/A response	FOR OFFICIAL USE ONLY
Height, Weight and Body Mass Index	Measurements for children. Ages: (0) to (11) months, (1) to (4) years, (5) to (10) years, (11) to (14) years, (15) to (17) years.		
Hematocrit or hemoglobin screening	Screening for all children		
Hemoglobinopathies screening: newborns	Screening for sickle cell disease in newborns.		
Hypothyroidism screening: newborns	Screening for congenital hypothyroidism in newborns.		
Immunization	Vaccines for children from birth to age (18), doses, recommended ages, and recommended populations vary: Diphtheria, Tetanus, Pertussis, Haemophilus influenza type b, Hepatitis A, Hepatitis B, Human Papillomavirus, Inactivated Poliovirus, Influenza (Flu Shot), Measles, Mumps, Rubella, Meningococcal, Pneumococcal, Rotavirus, Varicella. Catch ups must be covered.		
	HPV vaccination beginning at age (9) years for children and youth with any history of sexual abuse or assault who have not initiated or complete the (3) doses series (ACIP recommendation).		
	COVID-19 vaccine as part of preventive immunizations for adolescents over 16 years of age and adults, according to the order established by the Puerto Rico Department of Health and the recommendations of the Advisory Committee on Immunization Practices (ACIP), of the Centers for Disease Control and Prevention (CDC), in accordance with Article 2.050 (C) of the Health Insurance Code of Puerto Rico.		Upo
	Dengue vaccine for children from Age 9 to 16 years living in dengue endemic areas AND have laboratory confirmation of previous dengue infection - 3-dose series administered at 0, 6, and 12 months according to the recommendations of the Advisory Committee on Immunization Practices (ACIP) and the Centers for Disease Control and Prevention (CDC).		

<sup>\*</sup> Page number range are not accepted (i.e. 28-32)
\*\* The absence of explanation may prevent the filing approval.

<sup>&</sup>lt;sup>1</sup> Term use in this attachment to refer to an insured or subscriber

Benefit	Description	Please specify location (Form/Page/Paragraph/ Other) of complying provision/language or attach explanation for an N/A response	FOR OFFICIAL USE ONLY
Lead screening	For children at risk of lead exposure		
Maternal depression	Screening for mothers of infants at 1, 2, 4, and 6-month visits		
Medical History	For all children throughout development Ages: (0) to (21) years.		

<sup>\*</sup> Page number range are not accepted (i.e. 28-32)

\*\* The absence of explanation may prevent the filing approval.

<sup>&</sup>lt;sup>1</sup> Term use in this attachment to refer to an insured or subscriber

Benefit	Description	Please specify location (Form/Page/Paragraph/ Other) of complying provision/language or attach explanation for an N/A response	FOR OFFICIAL USE ONLY	
Obesity screening: children and adolescents	The USPSTF recommends that clinicians screen children age (6) years and older for obesity and offer them, or refer them to comprehensive, intensive behavioral interventions to promote improvement in weight status.			
Oral Health	Risk assessment for young children. Ages: (0) to (11) months, (1) to (4) years, (5) to (10) years.			
Phenylketonuria (PKU) screening: newborns	Screening for phenylketonuria in newborns.			
Prevention of Dental Caries in Children Younger Than 5 Years: Screening and Interventions: children younger than 5 years	The USPSTF recommends that primary care clinicians prescribe oral fluoride supplementation starting at age 6 months for children whose water supply is deficient in fluoride.  The USPSTF recommends that primary care clinicians apply fluoride varnish to the primary			Updated
	teeth of all infants and children starting at the age of primary tooth eruption.			
Skin Cancer behavioral counseling	The USPSTF recommends counseling young adults, adolescents, children, and parents of young children about minimizing exposure to ultraviolet (UV) radiation for persons aged 6 months to 24 years with fair skin types to reduce their risk of skin cancer.			
Tobacco use: children and adolescents	The USPSTF recommends that primary care clinicians provide interventions, including education or brief counseling, to prevent initiation of tobacco use among school-aged children and adolescents.			
Tobacco, Alcohol or Drug Use	Screening children (11) to (21) years of age.			
Tuberculosis	Testing for children at higher risk of tuberculosis. Ages: (0) to (11) months, (1) to (4) years, (5) to (10) years, (11) to (14) years, (15) to (21) years.			
Vision screening: children	The USPSTF recommends vision screening for all children at least once between the ages of (3) and (5) years, to detect the presence of amblyopia or its risk factors.			

<sup>\*</sup> Page number range are not accepted (i.e. 28-32)
\*\* The absence of explanation may prevent the filing approval.

<sup>&</sup>lt;sup>1</sup> Term use in this attachment to refer to an insured or subscriber

Benefit	Description	Please specify location (Form/Page/Paragraph/ Other) of complying provision/language or attach explanation for an N/A response	FOR OFFICIAL USE ONLY
---------	-------------	--	--------------------------

#### References

https://www.uspreventiveservicestaskforce.org/uspstf/recommendation-topics/uspstf-and-b-recommendations

 $\underline{https://mchb.hrsa.gov/programs-impact/programs/preventive-guidelines-screenings-women-children-youth}$ 

https://www.womenspreventivehealth.org/recommendations/

https://downloads.aap.org/AAP/PDF/periodicity\_schedule.pdf

https://www.cdc.gov/vaccines/schedules/downloads/adult/adult-combined-schedule.pdf

https://www.cdc.gov/vaccines/schedules/downloads/child/0-18yrs-child-combined-schedule.pdf

#### CERTIFICATION

I have reviewed	or supervised the preparation of the above checklist and certify that the forms(s) and/or dru	ig formulary is (are) in compliance with all of the
Signatu	ire:	
Date	:	

<sup>\*</sup> Page number range are not accepted (i.e. 28-32)

<sup>\*\*</sup> The absence of explanation may prevent the filing approval.

<sup>&</sup>lt;sup>1</sup> Term use in this attachment to refer to an insured or subscriber